Author Index

ABE K.Y. *see* KUO P.C.
ABE T. *see* KORENAGA M.
ABRAHAM-INPIJN L. *see* WASSENAAR A.
ACTOR J.K. *see* GIESE N.A.
ADAMS S.E. *see* LAYTON G.T.
AGIS H., FÜREDER W., BANKL H.C., KUNDI M., SPERR W.R., WILLHEIM M., BOLTZ-NITULESCU G., BUTTERFIELD J.H., KISHI K., LECHNER K. & VALENT P. Comparative immunophenotypic analysis of human mast cells, blood basophils and monocytes, 535
AKOUM H., TSICOPOULOS A., VORNG H., WALLAERT B., DESSAINT J.P., JOSEPH M., HAMID Q. & TONNEL A.B. Venom immunotheraphy modulates interleukin-4 and interferon-γ messenger RNA expression of peripheral T lymphocytes, 593
ALFREY E.J. *see* KUO P.C.
ALLING D.W. *see* DUCHMANN R.
AMOUI M *see* DRÁBEROVÁ L.
ANDERSON G. *see* TCHILIAN E.Z.
ARDESTANI S. *see* LIANG B.
AUDETTE M. *see* BOSSÉ M.
AXELSSON B. *see* YOUSEFFI-ETEMAD R.
AYROLDI E., CANNARILE L. & RICCARDI C. Modulation of superantigen-induced T-cell deletion by antibody anti-Pgp-1 (CD44), 191

BALDWIN C.L. *see* OKRAGLY A.J.
BALK S.P. *see* TAPLIN M.-E.
BANKL H.C. *see* AGIS H.
BARAK V. *see* LEVI-SCHAFFER F.
BARKANS J. *see* LEVI-SCHAFFER F.
BARNARD A., MAHON B.P., WATKINS J., REDHEAD K. & MILLS K.H.G. Th1/Th2 cell dichotomy in acquired immunity to *Bordetella pertussis*: variables in the *in vivo* priming and *in vitro* cytokine detection techniques affect the classification of T-cell subsets as Th1, Th2 or Th0, 372
BARNES P.J. *see* BERKMAN N.
BARNETSON R. StC. *see* CAVANAGH L.L.
BARRATT-BOYES S.M., HENDERSON R.A. & FINN O.J. Chimpanzee dendritic ells with potent immunostimulatory function can be propagated from peripheral blood, 528
BARRETT A.J. *see* JIANG Y.Z.
BASSETT H.F. *see* DOHERTY M.L.
BATTEN P., YACOUB M.H. & ROSE M.L. Effect of human cytokines (IFN-γ, TNF-α, IL-1β, IL-4) on porcine endothelial cells: induction of MHC and adhesion molecules and functional significance of these changes, 127
BECKHOVE P. *see* UMANSKY V.
BELANI R. & WEINER G.J. Expression of both B7-1 and CD28 contributes to the IL-2 responsiveness of CTLL-2 cells, 271
BERGER M.F. *see* SEWELL W.A.
BERKMAN N., ROBICHAUD A., KRISHNAN V.L., ROESEMS G., ROBBINS R., JOSE P.J., BARNES P.J. & CHUNG K.F. Expression of RANTES in human airway epithelial cells: effect of corticosteroids and interleukin-4, -10 and -13, 599
BERTRAND A. *see* JURCEVIC S.
BETTS C. *see* WILSON R.A.
BIANCHI A.T.J. *see* CUKROWSKA B.
BIERHAUS R. *see* BOEHME N.W.J.
BJÖRCK P. & PAULIE S. CD40 antibodies defining distinct epitopes display qualitative differences in their induction of B-cell differentiation, 291
BLAND P. *see* PARSONS K.R.
BLUMBERG R.S. *see* TAPLIN M.-E.
BODMAN-SMITH K.B. *see* JEDDI P.A.
BOEHME N.W.J., DENG Y., RAETH U., BIERHAUS R., ZIEGLER R., STREMMEL W. & NAWROTH P.P. Release of thrombomodulin from endothelial cells by concerted action of TNF-α and neutrophils: *in vivo* and *in vitro* studies, 134
BOLTZ-NITULESCU G. *see* AGIS H.
BOOG C.J.P. *see* WIJBURG O.L.C.
BOSHELL M., MCLEOD J., WALKER L., HALL N., PATEL Y. & SANSOM D. Effects of antigen presentation on superantigen-induced apoptosis mediated by Fas/Fas ligand interactions in human T cells, 586
BOSSÉ M., AUDETTE M., FERLAND C., PELLETIER G., CHU H.W., DAKHAMA A., LAVIGNE S., BOULET L.-P. & LAVIOLETTE M. Gene expression of interleukin-2 in purified human peripheral blood eosinophils, 149
BOST K.L., HOLTON R.H., KINCY CAIN T. & CLEMENTS J.D. *In vivo* treatment with anti-interleukin-13 antibodies significantly reduces the humoral immune response against an oral immunogen in mice, 633
BOULET L.-P. *see* BOSSÉ M.
BRACKERTZ D. *see* TRINDER P.K.E.
BRADBURY M.G., DOHERTY K.V., PARISH C.R. & LYONS A.B. The immunosuppressive compound 2-acetyl-4-tetrahydroxybutyl imidazole inhibits the allogeneic mixed lymphocyte reaction by sequestration of a recirculating subpopulation of T cells, 80
BRAUN R.K. *see* STERNER-KOCK A.
BRETT S.J. & TITE J.P. Both H-2- and non-H-2-linked genes influence influenza nucleoprotein epitope recognition by $CD4^+$ T cells, 42
BROSSARD M. *see* GANAPAMO F.
BUTTERFIELD J.H. *see* AGIS H.

CAI J.-P. *see* CHIN Y.-H.
CAMBRIDGE G. *see* LONGHURST C.
CAMPBELL D.G. *see* POLLOCK J.M.
CANNARILE L. *see* AYROLDI E.
CANNING C. *see* TAPLIN M.-E.
CARO H.N. *see* SHEIKH N.A.
CASTEELS-VAN DAELE M. *see* KRÜGER M.
CAVANAGH L.L., SLUYTER R., HENDERSON K.G., BARNETSON R. StC. & HALLIDAY G.M. Epidermal Langerhans' cell induction of immunity against an ultraviolet-induced skin tumour, 475
CEUPPENS J.L. *see* KRÜGER M.
CHAPMAN C. *see* LONGHURST C.
CHEN Z. *see* GORCZYNSKI R.M.
CHIN Y.-H., YE M.-W., CAI J.-P. & XU X.-M. Differential regulation of tissue-specific lymph node high endothelial venule cell adhesion molecules by tumour necrosis factor and transforming growth factor-β_1, 559
CHONG P. *see* ZHANG L.
CHRISTIANSEN D., MILLAND J., THORLEY B.R., MCKENZIE I.F.C., MOTTRAM P.L., PURCELL L.J. & LOVELAND B.E. Engineering of recombinant soluble CD46: an inhibitor of complement activation, 348
CHRISTMAS S.E. *see* DENIZ G.
see also GUDELJ L.
CHU H.W. *see* BOSSÉ M.
CHUNG H.T. *see* PARK Y.C.
CHUNG K F. *see* BERKMAN N.
CHUNG S. *see* GORCZYNSKI R.M.
CLEMENTS J.D. *see* BOST K.L.
COHEN S.B.A., WEBB L.M.C. & FELDMANN M. The method of deriving human T-cell clones alters the proportion of IL-10-producing cells, 343
COLE J.S. *see* LAYTON G.T.

COOLEY M.A. *see* SEWELL W.A.
COOREVITS L. *see* KRÜGER M.
COPPIN H.L. *see* JURCEVIC S.
COSTA M.G. *see* GIROLOMONI G.
COULSON A.F.W. *see* MUSTAFA A.S.
COULSON P.S. *see* WILSON R.A.
COURIEL D. *see* JIANG Y.Z.
CROCKARD A.D. *see* POLLOCK J.M.
CROCKER P.R. *see* UMANSKY V.
CUKROWSKA B., ŠINKORA J., MANDEL L., SPLÍCHAL I., BIANCHI A.T.J., KOVÁŘŮ F. & TLASKALOVÁ-HOGENOVÁ H. Thymic B cells of pig fetuses and germ-free pigs spontaneously produce IgM, IgG and IgA: detection by ELISPOT method, 487
CUMBERBATCH M., DEARMAN R.J. & KIMBER I. Constitutive and inducible expression of interleukin-6 by Langerhans' cells and lymph node dendritic cells, 513

DAFFERN P.J. *see* DISCIPIO R.G.
DAFOE D.C. *see* KUO P.C.
DAKHAMA A. *see* BOSSÉ M.
DANNECKER G.E. *see* SCHULTZ H.
DAVIS W.C. *see* DOHERTY M.L.
DE BAETSELIER P. *see* THORP K.M.
DE PREVAL C. *see* JURCEVIC S.
DE WIT T.P.M. *see* KRÜGER M.
DEARMAN R.J. *see* CUMBERBATCH M.
DELVES P.J. *see* JEDDI P.A.
DENG Y. *see* BOEHMÉ N.W.J.
DENIZ G. *see* GUDELJ L.
DENIZ G., CHRISTMAS S.E. & JOHNSON P.M. Soluble mediators and cytokines produced by human $CD3^-$ leucocyte clones from decidualized endometrium, 92
DERMIME S. *see* JIANG Y.Z.
DESSAINT J.P. *see* AKOUM H.
DINARELLO C.A. *see* ENDRES S.
DISCIPIO R.G., DAFFERN P.J., KAWAHARA M., PIKE R., TRAVIS J. & HUGLI T.E. Cleavage of human complement component C5 by cysteine proteinases from *Porphyromonas (Bacteriodes) gingivalis*. Prior oxidation of C5 augments proteinase digestion of C5, 660
DOCKRELL H.M. *see* TURNER J.
DOHERTY K.V. *see* BRADBURY M.G.
DOHERTY M.L., BASSETT H.F., QUINN P.J., DAVIS W.C., KELLY A.P. & MONAGHAN M.L. A sequential study of the bovine tuberculin reaction, 9
DOWLING M.-A. *see* WILSON R.A.
DRÁBER P *see* DRÁBEROVÁ L.
DRÁBEROVÁ L., AMOUI M & DRÁBER P Thy-1-mediated activation of rat mast cells: the role of Thy-1 membrane microdomains, 141
DUCHMANN R., JAFFE J., EHRHARDT R., ALLING D.W. & STROBER W. Differential usage of T-cell receptor $V\beta$ gene families by $CD4^+$ and $CD8^+$ T cells in patients with $CD8^{hi}$ common variable immunodeficiency: evidence of a post-thymic effect, 99

EHRENSTEIN M.R. *see* LONGHURST C.
EHRHARDT R. *see* DUCHMANN R.
ELSER C. *see* MUNK M.E.
ENDRES S., WHITAKER R.E.D., GHORBANI R., MEYDANI S.N. & DINARELLO C.A. Oral aspirin and ibuprofen increase cytokine-induced synthesis of IL-1β and of tumour necrosis factor-α *ex vivo*, 264
ERICKSON K.L. *see* SUK K.
ESHIMA K., SUZUKI H., YAMAZAKI S. & SHINOHARA N. Derivation of T-cell receptor α-chain double expresser lines from normal murine mature T cells, 205

FAINBOIM L. *see* LARREGINA A.
FAZLE AKBAR S.M., INABA K. & ONJI M. Upregulation of MHC class II antigen on dendritic cells from hepatitis B virus transgenic mice by interferon-γ: abrogation of immune response defect to a T-cell-dependent antigen, 519
FELDMANN M. *see* COHEN S.B.A.
FERLAND C. *see* BOSSÉ M.
FINN O.J. *see* BARRATT-BOYES S.M.
FRANTZ M.E. *see* TAPLIN M.-E.
FRENCH T.J. *see* LAYTON G.T.
FROST V. *see* KAY J.E.
FÜREDER W. *see* AGIS H.
FUSCO M. *see* ZOLA H.

GALLI S.J. *see* HILL P.B.
GANAPAMO F., RUTTI B. & BROSSARD M. Immunosuppression and cytokine production in mice infested with *Ixodes ricinus* ticks: a possible role of laminin and interleukin-10 on the *in vitro* responsiveness of lymphocytes to mitogens, 259
GASKINS H.R. *see* ZUCKERMANN F.A.
GAZZINELLI R.T. *see* GIESE N.A.
GEISELHART A. *see* SCHULTZ H.
GHORBANI R. *see* ENDRES S.
GIANNETTI A. *see* GIROLOMONI G.
GIENI R.S., YANG X., KELSO A. & HAYGLASS K.T. Limiting dilution analysis of CD4 T-cell cytokine production in mice administered native versus polymerized ovalbumin: directed induction of T-helper type-1-like activation, 119
GIESE N.A., GAZZINELLI R.T., ACTOR J.K., MORAWETZ R.A., SARZOTTI M. & MORSE H.C.III. Retrovirus-elicited interleukin-2 and tumour necrosis factor-α as inducers of interferon-γ-mediated pathology in mouse AIDS, 467
GIROLOMONI G., VALLE M.T., ZACCHI V., COSTA M.G., GIANNETTI A. & MANCA F. Cultured human Langerhans' cells are superior to fresh cells at presenting native HIV-1 protein antigens to specific $CD4^+$ T-cell lines, 310
GIRVIN R.M. *see* POLLOCK J.M.
GORCZYNSKI R.M., CHEN Z., HOANG Y. & ROSSI-BERGMAN B. A subset of $\gamma\delta$ T-cell receptor-positive cells produce T-helper type-2 cytokines and regulate mouse skin graft rejection following portal venous pretransplant preimmunization, 381
GORCZYNSKI R.M., CHUNG S., HOANG Y., SULLIVAN B. & CHEN Z. Altered patterns of migration of cytokine-producing T lymphocytes in skin-grafted naive or immune mice following *in vivo* administration of anti-VCAM-1 or -ICAM-1, 573
GOTCH F. *see* LAYTON G.T.
GRDIĆ D. *see* HÖRNQUIST E.
GUDELJ L., DENIZ G., RUKAVINA D., JOHNSON P.M. & CHRISTMAS S.E. Expression of functional molecules by human $CD3^-$ decidual granular leucocyte clones, 609

HALL N. *see* BOSHELL M.
HALLIDAY G.M. *see* CAVANAGH L.L.
HAMID Q. *see* AKOUM H.
HAN S.-H. *see* SUN K-H
HANBY-FLARIDA M. *see* OKRAGLY A.J.
HARADA M., OKAMOTO T., OMOTO K., TAMADA K., TAKENOYAMA M., HIRASHIMA C., ITO O., KIMURA G. & NOMOTO K. Specific immunotherapy with tumour-draining lymph node cells cultured with both anti-CD3 and anti-CD28 monoclonal antibodies, 447
HARRIS S.J. *see* LAYTON G.T.
HASHIGUCHI Y. *see* KORENAGA M.
HASSAN J. & REEN D.J. Reduced primary antigen-specific T-cell precursor frequencies in neonates is associated with deficient interleukin-2 production, 604
HAYGLASS K.T. *see* GIENI R.S.
HEEMSKERK M.H.M. *see* WIJBURG O.L.C.
HENDERSON K.G. *see* CAVANAGH L.L.
HENDERSON R.A. *see* BARRATT-BOYES S.M.
HENSEL N.F. *see* JIANG Y.Z.
HILL P.B., MACDONALD A.J., THORNTON E.M., NEWLANDS G.F.J., GALLI S.J. & MILLER H.R.P. Stem cell factor enhances immunoglobulin E-dependent mediator release from cultured rat bone marrow-derived mast cells: activation of previously unresponsive cells demonstrated by a novel ELISPOT assay, 326
HILL-PERKINS M. *see* LAYTON G.T.

HIRASHIMA C. *see* HARADA M.
HIROMATSU K. *see* KIMURA Y.
HOANG Y. *see* GORCZYNSKI R.M.
HOFFMANN M.K. *see* SCHULTZ H.
HOHENSTEIN R. *see* LEVI-SCHAFFER F.
HOLT P.G. *see* STRICKLAND D.
HOLTON R.H. *see* BOST K.L.
HÖRNQUIST E., GRDIĆ D., MAK T. & LYCKE N. CD8-deficient mice exhibit augmented mucosal immune responses and intact adjuvant effects to cholera toxin, 220
HOWARD C.J. *see* PARSONS K.R.
HSIEH S.-C. *see* SUN K-H
HUGLI T.E. *see* DISCIPIO R.G.
HUIE P. *see* KUO P.C.
HUSMANN R.J. *see* ZUCKERMANN F.A.
HYDE D.M. *see* STERNER-KOCK A.

IGIETSEME J.U. The molecular mechanism of T-cell control of *Chlamydia* in mice: role of nitric oxide, 1
IKUTA K. *see* KODAIRA Y.
INABA K. *see* FAZLE AKBAR S.M.
ISENBERG D.A. *see* JEDDI P.A.
see also LONGHURST C.
ITO O. *see* HARADA M.
IWAKAMI N., KIKUCHI A., KUNISHITA T., YAMAMOTO H., NONAKA I. & KAMO I. Analysis of lymphoproliferative cytokines produced by thymic myoid cells, 108

JAFFE J. *see* DUCHMANN R.
JEDDI P.A., BODMAN-SMITH K.B., LUND T., LYDYARD P.M., MENGLE-GAW L., ISENBERG D.A., YOUINOU P. & DELVES P.J. Agalactosyl IgG and β-1,4-galactosyltransferase gene expression in rheumatoid arthritis patients and in the arthritis-prone MRL *lpr/lpr* mouse, 654
JENKINSON E.J. *see* TCHILIAN E.Z.
JIANG Y.Z., COURIEL D., MAVROUDIS D.A., LEWALLE P., MALKOVSKA V., HENSEL N.F., DERMIME S., MOLLDREM J. & BARRETT A.J. Interaction of natural killer cells with MHC class II: reversal of HLA-DR1-mediated protection of K562 transfectant from natural killer cell-mediated cytolysis by brefeldin-A, 481
JOHNSON P.M. *see* DENIZ G.
see also GUDELJ L.
JONES B.V. *see* PARSONS K.R.
JOSE P.J. *see* BERKMAN N.
JOSEPH M. *see* AKOUM H.
JUN C.D. *see* PARK Y.C.
JURCEVIC S., PRAUD C., COPPIN H.L., BERTRAND A., RICARD S., THOMSEN M., LAKHDAR-GHAZAL F. & DE PREVAL C. Role of polymorphic residues of human leucocyte antigen-DR molecules on the binding of human immunodeficiency virus peptides, 414

KAMO I. *see* IWAKAMI N.
KANEGANE H., KASAHARA Y., NIIDA Y., YACHIE A., SUGII S., TAKATSU K., TANIGUCHI N. & MIYAWAKI T. Expression of L-selectin (CD62L) discriminates Th1- and Th2-like cytokine-producing memory $CD4^+$ T cells, 186
KANG H.S. *see* PARK Y.C.
KASAHARA Y. *see* KANEGANE H.
KAUFMANN S.H.E. *see* MUNK M.E.
KAWAHARA M. *see* DISCIPIO R.G.
KAY A.B. *see* LEVI-SCHAFFER F.
KAY J.E., SMITH M.C., FROST V. & MORGAN G.Y. Hypersensitivity to rapamycin of BJAB B lymphoblastoid cells, 390
KEES U.R. *see* STRICKLAND D.
KELLY A.P. *see* DOHERTY M.L.
KELLY C.G. *see* TODRYK S.M.
KELSO A. *see* GIENI R.S.
KIEVITS F. *see* WASSENAAR A.
KIKUCHI A. *see* IWAKAMI N.
KIM H.D. *see* PARK Y.C.
KIM H.M. *see* PARK Y.C.
KIMBER I. *see* CUMBERBATCH M.
KIMURA G. *see* HARADA M.
KIMURA Y., TOMIDA S., MATSUMOTO Y., HIROMATSU K. & YOSHIKAI Y. Evidence for the early recruitment of T-cell receptor $\gamma\delta^+$ T cells during rat listeriosis, 21
KINCY CAIN T. *see* BOST K.L.
KINGSMAN A.J. *see* LAYTON G.T.
KIRKHAM P.A., TAKAMATSU H., YANG H. & PARKHOUSE R.M.E. Porcine CD3ϵ: its characterization, expression and involvement in activation of porcine T lymphocytes, 616
KISHI K. *see* AGIS H.
KODAIRA Y., YOKOMURO K., TANAKA S., MIYAZAKI J.-I. & IKUTA K. Developmental heterogeneity of Vγ1.1 T cells in the mouse liver, 213
KOLKOWSKI E. *see* LARREGINA A.
KORENAGA M., WATANABE N., ABE T. & HASHIGUCHI Y. Acceleration of IgE responses by treatment with recombinant interleukin-3 prior to infection with *Trichinella spiralis* in mice, 642
KOVÁŘŮ F. *see* CUKROWSKA B.
KRIEGER N.R. *see* KUO P.C.
KRISHNAN V.L. *see* BERKMAN N.
KRÜGER A. *see* UMANSKY V.
KRÜGER M., COOREVITS L., DE WIT T.P.M., CASTEELS-VAN DAELE M., VAN DE WINKEL J.G.J. & CEUPPENS J.L. Granulocyte–macrophage colony-stimulating factor antagonizes the transforming growth factor-β-induced expression of FcγRIII (CD16) on human monocytes, 162
KUNDI M. *see* AGIS H.
KUNISHITA T. *see* IWAKAMI N.
KUO P.C., ALFREY E.J., KRIEGER N.R., ABE K.Y., HUIE P., SIBLEY R.K. & DAFOE D.C. Differential localization of allograft nitric oxide synthesis: comparison of liver and heart transplantation in the rat model, 647

LACY P. *see* LEVI-SCHAFFER F.
LAKHDAR-GHAZAL F. *see* JURCEVIC S.
LARREGINA A., MORELLI A., KOLKOWSKI E. & FAINBOIM L. Flow cytometric analysis of cytokine receptors on human Langerhans' cells. Changes observed after short-term culture, 317
LATCHMEN D. *see* LONGHURST C.
LAURENCE J. *see* MITRA D.
LAVIGNE S. *see* BOSSÉ M.
LAVIOLETTE M. *see* BOSSÉ M.
LAYTON G.T., HARRIS S.J., MYHAN J., WEST D., GOTCH F., HILL-PERKINS M., COLE J.S., MEYERS N., WOODROW S., FRENCH T.J., ADAMS S.E. & KINGSMAN A.J. Induction of single and dual cytotoxic T-lymphocyte responses to viral proteins in mice using recombinant hybrid Ty–virus-like particles, 171
LEAKER B. *see* LONGHURST C.
LECHNER K. *see* AGIS H.
LEHNER T. *see* TODRYK S.M.
LEVI-SCHAFFER F., BARKANS J., NEWMAN T.M., YING S., WAKELIN M., HOHENSTEIN R., BARAK V., LACY P., KAY A.B. & MOQBEL R. Identification of interleukin-2 in human peripheral blood eosinophils, 155
LEWALLE P. *see* JIANG Y.Z.
LIANG B., ARDESTANI S., MARCHALONIS J.J. & WATSON R.R. T-cell-receptor dose and the time of treatment during murine retrovirus infection for maintenance of immune function, 198
LIU W.-T. *see* SUN K-H
LONGHURST C., EHRENSTEIN M.R., LEAKER B., STEVENSON F.K., SPELLERBERG M., CHAPMAN C., LATCHMEN D., ISENBERG D.A. & CAMBRIDGE G. Analysis of immunoglobulin variable region genes of a human IgM anti-myeloperoxidase antibody derived from a patient with vasculitis, 334
LOOS M. *see* TRINDER P.K.E.
LOVELAND B.E. *see* CHRISTIANSEN D.
LUND T. *see* JEDDI P.A.
LUNDIN K.E.A. *see* MUSTAFA A.S.
LYCKE N. *see* HÖRNQUIST E.
LYDYARD P.M. *see* JEDDI P.A.

LYNCH D.H. *see* MITRA D.
LYONS A.B. *see* BRADBURY M.G.

MACARDLE P.J. *see* ZOLA H.
MACDONALD A.J. *see* HILL P.B.
MACKIE D.P. *see* POLLOCK J.M.
MAEURER M.J. *see* TRINDER P.K.E.
MAHON B.P. *see* BARNARD A.
MAK T. *see* HÖRNQUIST E.
MALKOVSKA V. *see* JIANG Y.Z.
MANCA F. *see* GIROLOMONI G.
MANDEL L. *see* CUKROWSKA B.
MANN D. *see* OKRAGLY A.J.
MARCHALONIS J.J. *see* LIANG B.
MARSHALL-CLARKE S., OWEN G. & TASKER L. Ligation of CD40 with soluble CD40 ligand reverses anti-immunoglobulin-mediated negative signalling in murine B lymphoma cell lines but not in immature B cells from neonatal mice, 624
MATSUMOTO Y. *see* KIMURA Y.
MATSUZAKI G., SONG F. & NOMOTO K. Suppression of T-helper type-1 immune response against *Listeria monocytogenes* by treatment of mice with goat antibodies to mouse IgD, 15
MATTHEWS N. *see* THORP K.M.
MAVROUDIS D.A. *see* JIANG Y.Z.
MCKENZIE I.F.C. *see* CHRISTIANSEN D.
MCLEOD J. *see* BOSHELL M.
MELOEN R.H. *see* MUSTAFA A.S.
MENGLE-GAW L. *see* JEDDI P.A.
MEYDANI S.N. *see* ENDRES S.
MEYERS N. *see* LAYTON G.T.
MILLAND J. *see* CHRISTIANSEN D.
MILLER H.R.P. *see* HILL P.B.
MILLS K.H.G. *see* BARNARD A.
MITRA D., STEINER M., LYNCH D.H., STAIANO-COICO L. & LAURENCE J. HIV-1 upregulates Fas ligand expression in $CD4^+$ T cells *in vitro* and *in vivo*: association with Fas-mediated apoptosis and modulation by aurintricarboxylic acid, 581
MIYAWAKI T. *see* KANEGANE H.
MIYAZAKI J.-I. *see* KODAIRA Y.
MOHAPATRA S.S. *see* ZHANG L.
MOLLDREM J. *see* JIANG Y.Z.
MONAGHAN M.L. *see* DOHERTY M.L.
MOORE N.C. *see* TCHILIAN E.Z.
MOQBEL R. *see* LEVI-SCHAFFER F.
MORAWETZ R.A. *see* GIESE N.A.
MORELLI A. *see* LARREGINA A.
MORGAN G.Y. *see* KAY J.E.
MORSE H.C. III. *see* GIESE N.A.
MOTTRAM P.L. *see* CHRISTIANSEN D.
MUNK M.E., ELSER C. & KAUFMANN S.H.E. Human γ/δ T-cell response to *Listeria monocytogenes* protein components *in vitro*, 230
MUNRO G.H. *see* TODRYK S.M.
MUSTAFA A.S., LUNDIN K.E.A., MELOEN R.H., SHINNICK T.M., COULSON A.F.W. & OFTUNG F. HLA-DR4-restricted T-cell epitopes from the microbacterial 60 000 MW heat shock protein (hsp 60) do not map to the sequence homology regions with the human hsp 60, 414
MYHAN J. *see* LAYTON G.T.

NAWROTH P.P. *see* BOEHME N.W.J.
NEILL S.D. *see* POLLOCK J.M.
NEWLANDS G.F.J. *see* HILL P.B.
NEWMAN I. & WILKINSON P.C. The bacterial superantigen Staphylococcal enterotoxin B stimulates lymphocyte locomotor capacity during culture *in vitro*, 428
NEWMAN T.M. *see* LEVI-SCHAFFER F.
NIETHAMMER D. *see* SCHULTZ H.
NIIDA Y. *see* KANEGANE H.
NOMOTO K. *see* HARADA M.
see also MATSUZAKI G.
NONAKA I. *see* IWAKAMI N.

OFTUNG F. *see* MUSTAFA A.S.
OHSUGI K. *see* SUZUKI Y.
OKAMOTO T. *see* HARADA M.
OKRAGLY A.J., HANBY-FLARIDA M., MANN D. & BALDWIN C.L. Bovine γ/δ T-cell proliferation is associated with self-derived molecules constitutively expressed *in vivo* on mononuclear phagocytes, 71
OMOTO K. *see* HARADA M.
ONJI M. *see* FAZLE AKBAR S.M.
ONO Y. *see* SUZUKI Y.
OWEN G. *see* MARSHALL-CLARKE S.
OWEN J.J.T. *see* TCHILIAN E.Z.

PARISH C.R. *see* BRADBURY M.G.
PARK Y.C., JUN C.D., KANG H.S., KIM H.D., KIM H.M. & CHUNG H.T. Role of intracellular calcium as a priming signal for the induction of nitric oxide synthesis in murine peritoneal macrophages, 296
PARKHOUSE R.M.E. *see* KIRKHAM P.A.
PARSONS K.R., SOPP P., JONES B.V., BLAND P. & HOWARD C.J. Identification of a molecule uniquely expressed on a γ/δ TCR^+ subset within bovine intestinal intraepithelial lymphocytes, 64
PARTIDOS C.D., VOHRA P. & STEWARD M.W. Induction of measles virus-specific cytotoxic T-cell responses after intranasal immunization with synthetic peptides, 179
PATEL Y. *see* BOSHELL M.
PAULIE S. *see* BJÖRCK P.
PELLETIER G. *see* BOSSÉ M.
PIKE R. *see* DISCIPIO R.G.
PLAYFAIR J.H.L. *see* SHEIKH N.A.
POLLOCK D.A. *see* POLLOCK J.M.
POLLOCK J.M., POLLOCK D.A., CAMPBELL D.G., GIRVIN R.M., CROCKARD A.D., NEILL S.D. & MACKIE D.P. Dynamic changes in circulating and antigen-responsive T-cell subpopulations post-*Mycobacterium bovis* infection in cattle, 236
PRAUD C. *see* JURCEVIC S.
PURCELL L.J. *see* CHRISTIANSEN D.

QUINN P.J. *see* DOHERTY M.L.

RADEMACHER T.W. *see* SHEIKH N.A.
RAETH U. *see* BOEHME N.W.J.
RAKMILEVICH L. *see* SAYLES P.C.
REDHEAD K. *see* BARNARD A.
REEN D.J. *see* HASSAN J.
REINHARDUS T. *see* WASSENAAR A.
RICARD S. *see* JURCEVIC S.
RICCARDI C. *see* AYROLDI E.
RIDINGS J. *see* ZOLA H.
RITZ J. *see* TAPLIN M.-E.
ROBBINS R. *see* BERKMAN N.
ROBERTON D.M. *see* ZOLA H.
ROBICHAUD A. *see* BERKMAN N.
ROCHA M. *see* UMANSKY V.
ROESEMS G. *see* BERKMAN N.
ROSE M.L. *see* BATTEN P.
ROSSI-BERGMAN B. *see* GORCZYNSKI R.M.
RUKAVINA D. *see* GUDELJ L.
RUTTI B. *see* GANAPAMO F.

SANDERS A. *see* WIJBURG O.L.C.
SANSOM D. *see* BOSHELL M.
SAPPLER G. *see* SCHULTZ H.
SARZOTTI M. *see* GIESE N.A.
SAYLES P.C. & RAKMILEVICH L. Exacerbation of *Plasmodium chabaudi* malaria in mice by depletion of TCR $\alpha\beta^+$ T cells, but not TCR $\gamma\delta^+$ T cells, 29
SCHIRRMACHER V. *see* UMANSKY V.
SCHRENZEL M.D. *see* STERNER-KOCK A.
SCHULTZ H., GEISELHART A., SAPPLER G., NIETHAMMER D., HOFFMANN M.K. & DANNECKER G.E. The superantigen *Staphylococcus* enterotoxin B induces a strong and accelerated secondary T-cell response rather than anergy, 49

SEWELL W.A., BERGER M.F., SKIPSEY L.J., COOLEY M.A. & WARREN H.S. Cytokine expression by high-density human lymphocytes, 408
SHEIKH N.A., CARO H.N., TAVERNE J., PLAYFAIR J.H.L. & RADEMACHER T.W. Malaria: a tumour necrosis factor inhibitor from parasitized erythrocytes, 461
SHINNICK T.M. *see* MUSTAFA A.S.
SHINOHARA N. *see* ESHIMA K.
SIBLEY R.K. *see* KUO P.C.
ŠINKORA J. *see* CUKROWSKA B.
SKIPSEY L.J. *see* SEWELL W.A.
SLUYTER R. *see* CAVANAGH L.L.
SMITH M.C. *see* KAY J.E.
SMYTHIES L.E. *see* WILSON R.A.
SONG F. *see* MATSUZAKI G.
SOPP P. *see* PARSONS K.R.
SOUTHERN C. *see* THORP K.M.
SPECTER S. *see* ZHENG Z.M.
SPELLERBERG M. *see* LONGHURST C.
SPERR W.R. *see* AGIS H.
SPLÍCHAL I. *see* CUKROWSKA B.
STAIANO-COICO L. *see* MITRA D.
STEINER M. *see* MITRA D.
STERNER-KOCK A., BRAUN R.K., SCHRENZEL M.D. & HYDE D.M. Recombinant tumour necrosis factor-α and platelet-activating factor synergistically increase intercellular adhesion molecule-1 and E-selectin-dependent neutrophil adherence to endothelium *in vitro*, 454
STEVENSON F.K. *see* LONGHURST C.
STEWARD M.W. *see* PARTIDOS C.D.
STREMMEL W. *see* BOEHME N.W.J.
STRICKLAND D., KEES U.R. & HOLT P.G. Regulation of T-cell activation in the lung: alveolar macrophages induce reversible T-cell anergy *in vitro* associated with inhibition of interleukin-2 receptor signal transduction, 250
STRICKLAND D., KEES U.R. & HOLT P.G. Regulation of T-cell activation in the lung: isolated lung T cells exhibit surface phenotypic characteristics of recent activation including down-modulated T-cell receptors, but are locked into the G_0/G_1 phase of the cell cycle, 242
STROBER W. *see* DUCHMANN R.
SUGII S. *see* KANEGANE H.
SUK K. & ERICKSON K.L. Differential regulation of tumour necrosis factor-α mRNA degradation in macrophages by interleukin-4 and interferon-γ, 551
SULLIVAN B. *see* GORCZYNSKI R.M.
SUN K-H, LIU W.-T., TANG S.-J., TSAI C.-Y., HSIEH S.-C., WU T.-H., HAN S.-H. & YU C.-L. The expression of acidic ribosomal phosphoproteins on the surface membrane of different tissues in autoimmune and normal mice which are the target molecules for anti-double-stranded DNA antibodies, 362
SUZUKI H. *see* ESHIMA K.
SUZUKI Y., OHSUGI K. & ONO Y. Oxidative stress triggers tyrosine phosphorylation in B cells through a redox- and inflammatory cytokine-sensitive mechanism, 396

TAKAMATSU H. *see* KIRKHAM P.A.
TAKATSU K. *see* KANEGANE H.
TAKENOYAMA M. *see* HARADA M.
TAMADA K. *see* HARADA M.
TANAKA S. *see* KODAIRA Y.
TANG S.-J. *see* SUN K-H
TANIGUCHI N. *see* KANEGANE H.
TAPLIN M.-E., FRANTZ M.E., CANNING C., RITZ J., BLUMBERG R.S. & BALK S.P. Evidence against T-cell development in the adult human intestinal mucosa based upon lack of terminal deoxynucleotidyltransferase expression, 402
TASKER L. *see* MARSHALL-CLARKE S.
TAVERNE J. *see* SHEIKH N.A.
TCHILIAN E.Z., ANDERSON G., MOORE N.C., OWEN J.J.T. & JENKINSON E.J. Involvement of LFA-1/ICAM-2 adhesive interactions and PKC in activation-induced cell death following SEB rechallenge, 566
THOMSEN M. *see* JURCEVIC S.
THORLEY B.R. *see* CHRISTIANSEN D.
THORNTON E.M. *see* HILL P.B.
THORP K.M., VERSCHUEREN H., DE BAETSELIER P., SOUTHERN C. & MATTHEWS N. Protein kinase C isotype expression and regulation of lymphoid cell motility, 434
TITE J.P. *see* BRETT S.J.
TLASKALOVÁ-HOGENOVÁ H. *see* CUKROWSKA B.
TODRYK S.M., KELLY C.G., MUNRO G.H. & LEHNER T. Induction of immune responses to functional determinants of a cell surface streptococcal antigen, 55
TOMIDA S. *see* KIMURA Y.
TONNEL A.B. *see* AKOUM H.
TRAVIS J. *see* DISCIPIO R.G.
TRINDER P.K.E., MAEURER M.J., BRACKERTZ D. & LOOS M. The collagen-like component of the complement system, Clq, is recognized by 7 S autoantibodies and is functionally impaired in synovial fluids of patients with rheumatoid arthritis, 355
TSAI C.-Y. *see* SUN K-H
TSICOPOULOS A. *see* AKOUM H.
TURNER J. & DOCKRELL H.M. Stimulation of human peripheral blood mononuclear cells with live *Mycobacterium bovis* BCG activates cytolytic $CD8^+$ T cells *in vitro*, 339

UMANSKY V., BECKHOVE P., ROCHA M., KRÜGER A., CROCKER P.R. & SCHIRRMACHER V. A role for sialoadhesin-positive tissue macrophages in host resistance to lymphoma metastasis *in vivo*, 303

VALENT P. *see* AGIS H.
VALLE M.T. *see* GIROLOMONI G.
VAN DE WINKEL J.G.J. *see* KRÜGER M.
VAN ROOIJEN N. *see* WIJBURG O.L.C.
VERSCHUEREN H. *see* THORP K.M.
VOHRA P. *see* PARTIDOS C.D.
VORNG H. *see* AKOUM H.

WAKELIN M. *see* LEVI-SCHAFFER F.
WALKER L. *see* BOSHELL M.
WALLAERT B. *see* AKOUM H.
WARREN H.S. *see* SEWELL W.A.
WASSENAAR A., REINHARDUS T., ABRAHAM-INPIJN L. & KIEVITS F. Type-1 and type-2 $CD8^+$ T-cell subsets isolated from chronic adult periodontitis tissue differ in surface phenotype and biological functions, 113
WATANABE N. *see* KORENAGA M.
WATKINS J. *see* BARNARD A.
WATSON R.R. *see* LIANG B.
WEBB L.M.C. *see* COHEN S.B.A.
WEEDON H. *see* ZOLA H.
WEINER G.J. *see* BELANI R.
WEST D. *see* LAYTON G.T.
WHITAKER R.E.D. *see* ENDRES S.
WIJBURG O.L.C., HEEMSKERK M.H.M., SANDERS A., BOOG C.J.P. & VAN ROOIJEN N. Role of virus-specific $CD4^+$ cytotoxic T cells in recovery from Mouse Hepatitis Virus infection, 34
WILKINSON P.C. *see* NEWMAN I.
WILLHEIM M. *see* AGIS H.
WILSON R.A., COULSON P.S., BETTS C., DOWLING M.-A. & SMYTHIES L.E. Impaired immunity and altered pulmonary responses in mice with a disrupted interferon-γ receptor gene exposed to the irradiated *Schistosoma mansoni* vaccine, 275
WOODROW S. *see* LAYTON G.T.
WU T.-H. *see* SUN K-H

XU X.-M. *see* CHIN Y.-H.

YACHIE A. *see* KANEGANE H.
YACOUB M.H. *see* BATTEN P.
YAMAMOTO H. *see* IWAKAMI N.
YAMAZAKI S. *see* ESHIMA K.
YANG H. *see* KIRKHAM P.A.
YANG M. *see* ZHANG L.
YANG X. *see* GIENI R.S.

YE M.-W. *see* CHIN Y.-H.
YING S. *see* LEVI-SCHAFFER F.
YOKOMURO K. *see* KODAIRA Y.
YOSHIKAI Y. *see* KIMURA Y.
YOUINOU P. *see* JEDDI P.A.
YOUSEFFI-ETEMAD R. & AXELSSON B. Parallel pattern of expression of CD43 and LFA-1 on the $CD45RA^+$ (naive) and $CD45RO^+$ (memory) subsets of human $CD4^+$ and $CD8^+$ cells. Correlation with the aggregative response of the cells to CD43 monoclonal antibodies, 439
YU C.-L. *see* SUN K-H

ZACCHI V. *see* GIROLOMONI G.
ZHANG L., YANG M., CHONG P. & MOHAPATRA S.S. Multiple B- and T-cell epitopes on a major allergen of Kentucky Bluegrass pollen, 283
ZHENG Z.M. & SPECTER S. Dynamic production of tumour necrosis factor-α (TNF-α) messenger RNA, intracellular and extracellular TNF-α by murine macrophages and possible association with protein tyrosine phosphorylation of STAT1α and ERK2 as an early signal, 544
ZIEGLER R. *see* BOEHME N.W.J.
ZOLA H., FUSCO M., WEEDON H., MACARDLE P.J., RIDINGS J. & ROBERTON D.M. Reduced expression of the interleukin-2-receptor γ chain on cord blood lymphocytes: relationship to functional immaturity of the neonatal immune response, 86
ZUCKERMANN F.A. & GASKINS H.R. Distribution of porcine CD4/CD8 double-positive T lymphocytes in mucosa-associated lymphoid tissues, 493
ZUCKERMANN F.A. & HUSMANN R.J. Functional and phenotypic analysis of porcine peripheral blood CD4/CD8 double-positive T cells, 500

Subject Index

Adhesion molecules, high endothelial venule cell:
differential regulation by tumour necrosis factor and transforming growth factor β_1, 559
Agalactosyl IgG and galactosyltransferase gene expression:
in rheumatoid arthritis, 654
Allogeneic mixed lymphocyte reaction:
suppression by tetrahydroxybutyl imidazole, 80
Apoptosis, Fas mediated:
role of human immunodeficiency virus-1, 581
Apoptosis, Staphylococcal entertoxin B induced:
role of LFA-1/ICAM2 and PKC, 566
Apoptosis, superantigen induced:
mediated by Fas ligand in T cells, 586

B cells, thymic, porcine:
immunoglobulin production, 487
B-cell differentiation:
anti-CD40 antibodies and, 291

C1q, recognition by autoantibodies:
functional activity in rheumatoidarthritis, 355
CD3ε, porcine:
characterization, expression and role in T-lymphocyte activation, 616
$CD4^+$ cytotoxic T cells:
role in recovery from mouse hepatitis virus infection, 34
CD4/CD8 double-positive lymphocytes:
porcine, 493, 500
CD8-deficient mice:
augmented mucosal responses to Cholera toxin, 220
$CD8^+$ cytolytic T cells:
stimulation by *Mycobacterium bovis* bacillus Calmette–Guérin, 339
CD40 Ligand:
effect on growth inhibition of immature B cells, 624
CD43 and LFA-1:
expression on human T cells, 439
CD46, recombinant:
soluble, 348
Complement component C5, human:
cleavage by proteinases, 660
Cytokine expression:
by high-density lymphocytes, 408
Cytokine production, $CD4^+$:
induction of T-helper type-1-like activation, 119
Cytokine profile:
T-cell clones, 343
Cytokines, human:
effects on porcine endothelial cells, 127
Cytokines:
and cyclo-oxygenase inhibitors, 264
produced by thymic myoid cells, 108
Cytotoxic T lymphocyte responses, to viral proteins:
using virus-like particles, 171

Decidual leucocyte clones, $CD3^-$:
cytokine production, 92
Decidual leucocytes, $CD3^-$:
expression of function molecules, 609
Dendritic cells, chimpanzee:
propagation from peripheral blood, 528

Epitopes, B and T cell:
of Kentucky Bluegrass pollen, 283

Granulocyte–macrophage colony-stimulating factor:
inhibition of CD16 on human monocytes, 162

HLA-DR4-restricted T-cell epitopes:
of mycobacterial hsp60, 421
Human leucocyte antigen-DR molecules:
and human immunodeficiency virus peptide binding, 414

Immunotherapy, venom:
modulation of interleukin-4 and interferon-γ, 593
Immunotherapy:
with anti-CD28, 447
Influenza nucleoprotein:
epitope recognition, 42
Interferon-γ receptor gene:
disrupted and responses to *Schistosoma mansoni*, 275
Interleukin-2 responsiveness:
role of B7-1 and CD28 expression, 271
Interleukin-2-receptor γ chain:
on cord blood lymphocytes, 86
Interleukin-2:
detection in blood eosinophils, 149
in human eosinophils, 155
Interleukin-3:
acceleration of IgE responses, 642
role in humoral response to oral immunogen, 633
Interleukin-6 expression:
by Langerhans'and dendritic cells, 513
Intraepithelial lymphocytes, bovine:
antigen on γ/δ T-cell receptor$^+$ subset, 64
Intranasal immunization:
induction of cytotoxic T lymphocyte, 179

L-selectin:
and cytokine production by memory CD4 cells, 186
Laminin and interleukin-10:
and response to tick infestation, 259
Langerhans' cell, epidermal:
induction of immunity against ultraviolet-induced skin tumour, 475
Langerhans' cells, human:
cytokine receptors, 317
Langerhans' cells:
presentation of human immunodeficiency virus antigens, 310
Locomotion, lymphocyte:
stimulation by Staphylococcal enterotoxin B, 428

Major histocompatibility complex class II, upregulation on dendritic cells:
by interferon-γ, 519
Malaria, murine:
exacerbation in T-cell-depleted mice, 29
Mast cells, basophils, monocytes, human:
phenotypic analysis, 535
Motility, lymphocyte:
and protein kinase C isotype, 434
Mouse acquired immunodeficiency syndrome:
role of interleukin-12 and tumour necrosis factor-α in interferon-γ-induced pathology, 467

Natural killer cells:
interaction with major histocompatibility complex class II, 481
Neutrophil-endothelial adherence:
effect of tumour necrosis factor-α platelet-activating factor, 454
Nitric oxide synthesis, macrophage:
role of calcium, 296
Nitric oxide synthesis:
allograft, 647

Nitric oxide:
role in T-cell control of *Chlamydia*, 1

RANTES:
expression in airway epithelial cells, 599
Rapamycin:
hypersensitivity of B lymphoblastoid cells to, 390
Retrovirus infection, murine:
effects of T-cell receptor peptide immunization, 198
Ribosomal phosphoproteins:
targets for anti-double-stranded DNA antibodies, 362

Sialoadhesin-expressing macrophages:
in lymphoma, 303
Stem cell factor:
enhancement of IgE-dependent mediator release from mast cells, 326
Streptococcal antigen SA 1/11:
immune responses to, 55
Superantigen, induction of T-cell deletion:
modulation by anti-CD4, 191
Superantigen, *Staphylococcus* enterotoxin B:
accelerated T-cell response to, 49

T cells, cytokine producing:
altered patterns of migration in graft-draining nodes, 573
T-cell activation, lung:
anergy induced by lung macrophages, 250
T-cell activation, lung:
phenotypic analysis, 242
T-cell development:
in adult human intestinal mucosa, 402
T-cell precursor frequencies:
in newborns, 604
T-cell receptor α:
double-chain expresser lines, 205
T-cell receptor $\gamma\delta^+$ T cells:
during listeriosis, 21
T-cell receptor Vβ gene family usage:
in common variable immunodeficiency, 99
T-cell subpopulations, cattle:
to *Mycobacterium bovis*, 236
T-cell subsets, $CD8^+$:
in periodontitis, 113
T-cell subsets:
in acquired immunity to *Bordetella pertussis*, 372
$\gamma\delta$ T cells:
stimulation by self-derived molecules, 71
$\gamma\delta$ T-cell receptor T cells:
regulation of allograft rejection, 381
$\gamma\delta$ T-cell responses, human:
to *Listeria monocytogenes*, 230
T-helper type-1 response, to *Listeria*:
suppression by anti-IgD, 15
Thrombomodulin, release from endothelial cells:
by tumour necrosis factor-α and neutrophils, 134
Thy-1:
activation of mast cells, 141
Tuberculin reaction, bovine:
sequential study of, 9
Tumour necrosis factor inhibitor:
from malaria-parasitized erythrocytes, 461
Tumour necrosis factor-α messenger RNA, degradation:
regulation by interleukin-4 and interferon-γ, 551
Tumour necrosis factor-α:
and protein tyrosine phosphorylation, 544
Tyrosine phosphorylation, B cell:
triggering by oxidative stress, 396

Vγ1.1 T cells:
heterogeneity, 213
Variable region genes:
of human IgM anti-myeloperoxidase, 334